



---

Original article

# Social inequalities and COVID-19 mortality in the city of São Paulo, Brazil

Karina Braga Ribeiro,<sup>1\*</sup> Ana Freitas Ribeiro,<sup>2,3</sup>  
Maria Amélia de Sousa Mascena Veras<sup>1</sup> and Marcia Caldas de Castro<sup>4</sup>

<sup>1</sup>Faculdade de Ciências Médicas da Santa Casa de São Paulo, Department of Collective Health, São Paulo, Brazil, <sup>2</sup>Universidade Nove de Julho, São Paulo, Brazil, <sup>3</sup>Universidade Municipal de São Caetano do Sul, School of Medicine, São Caetano do Sul, Brazil and <sup>4</sup>Harvard TH Chan School of Public Health, Department of Global Health and Population, Boston, United States of America

\*Corresponding author. Faculdade de Ciências Médicas da Santa Casa de São Paulo, Department of Collective Health, Rua Doutor Cesário Motta Junior, 61–6o andar, São Paulo—SP Brazil, CEP 01221-020. E-mail: kbribeiro@gmail.com

Editorial decision 18 January 2021; Accepted 1 February 2021

## Abstract

**Background:** Heterogeneity in COVID-19 morbidity and mortality is often associated with a country's health-services structure and social inequality. This study aimed to characterize social inequalities in COVID-19 mortality in São Paulo, the most populous city in Brazil and Latin America.

**Methods:** We conducted a population-based study, including COVID-19 deaths among São Paulo residents from March to September 2020. Age-standardized mortality rates and unadjusted rate ratios (RRs) [with corresponding 95% confidence intervals (CIs)] were estimated by race, sex, age group, district of residence, household crowding, educational attainment, income level and percentage of households in subnormal areas in each district. Time trends in mortality were assessed using the Joinpoint model.

**Results:** Males presented an 84% increase in COVID-19 mortality compared with females (RR = 1.84, 95% CI 1.79–1.90). Higher mortality rates were observed for Blacks (RR = 1.77, 95% CI 1.67–1.88) and mixed (RR = 1.42, 95% CI 1.37–1.47) compared with Whites, whereas lower mortality was noted for Asians (RR = 0.63, 95% CI 0.58–0.68). A positive gradient was found for all socio-economic indicators, i.e. increases in disparities denoted by less education, more household crowding, lower income and a higher concentration of subnormal areas were associated with higher mortality rates. A decrease in mortality over time was observed in all racial groups, but it started earlier among Whites and Asians.

**Conclusion:** Our results reveal striking social inequalities in COVID-19 mortality in São Paulo, exposing structural inequities in Brazilian society that were not addressed by the governmental response to COVID-19. Without an equitable response, COVID-19 will further exacerbate current social inequalities in São Paulo.

**Key words:** COVID-19, mortality, social determinants of health, race factors

---

### Key Messages

- Males presented an 84% increase in COVID-19 mortality compared with females [relative risk (RR) = 1.84, 95% CI 1.79–1.90]. Higher mortality rates were observed for Blacks (RR = 1.77, 95% CI 1.67–1.88) and mixed (RR = 1.42, 95% CI 1.37–1.47) compared with Whites, whereas lower mortality was noted for Asians (RR = 0.63, 95% CI 0.58–0.68).
- A positive gradient was found for all socio-economic indicators, i.e. increases in disparities denoted by less education, more household crowding, lower income and a higher concentration of subnormal areas were associated with higher mortality rates.
- A decrease in mortality over time was observed in all racial groups, but it started earlier among Whites and Asians.
- Our finding of a higher percentage of deaths occurring in primary-care or isolated emergency-care units observed for Black and mixed populations may be related to barriers in healthcare access.
- Our results reveal striking social inequalities in COVID-19 mortality in São Paulo, exposing structural inequities in Brazilian society that were not addressed by the governmental response to COVID-19. Without an equitable response, COVID-19 will further exacerbate current social inequalities in São Paulo.

## Introduction

In December 2019, a cluster of pneumonia of unknown cause was registered in Wuhan, China and its etiologic agent, a new coronavirus, was named by the World Health Organization (WHO) as SARS-CoV-2 and the disease as COVID-19.<sup>1</sup> Given the spread of the disease worldwide, a pandemic was declared on 11 March.<sup>2</sup> According to WHO data, as of 12 October 2020, Brazil was the country with the third-highest number of cases, with 5 082 637 registered cases and 150 198 deaths.<sup>3</sup> The city of São Paulo is the most populous in the country (12.2 million inhabitants) and the first to report a COVID-19 case in Brazil. As of 14 October 2020, 302 223 cases and 13 196 deaths had been confirmed in the city, corresponding to crude incidence and mortality rates of 2467 and 108 per 100 000 inhabitants, respectively.<sup>4</sup>

Health inequities are recognized worldwide, with most vulnerable populations presenting higher incidence and mortality for communicable and non-communicable diseases. During the COVID-19 pandemic, socio-economic gradients were reported in Brazil,<sup>5,6</sup> the USA<sup>7–9</sup> and the UK.<sup>10</sup> Racial and ethnic disparities in infection and disease outcomes were described for many countries, with Blacks, Asians and minority ethnic groups emerging as more vulnerable than the White population.<sup>5,7–9,11</sup>

Heterogeneity in high rates of COVID-19 morbidity and mortality is often associated with a country's health structure and social inequality. Brazil is a highly unequal country, with substantial health and socio-economic differences observed between and within cities.<sup>12</sup> Previous studies have shown racial and socio-economic differences on COVID-19 mortality in Brazil,<sup>5,6</sup> but there are still gaps in knowledge regarding the interplay between race and socio-economic factors. This study addresses this issue and aims

to characterize mortality patterns from COVID-19 in the city of São Paulo, Brazil and to identify risk factors of mortality considering various socio-demographic indicators.

## Methods

### Study design: population-based descriptive study

#### Data source

Data on COVID-19 mortality among residents in the city of São Paulo were extracted from the Mortality Information System.<sup>13</sup> These include all deaths classified with a confirmed code B34.2 or a suspected code U04.9, based on the International Classification of Diseases version 10 (ICD-10). We included deaths that occurred between March and September 2020. Each record had information on age (<4, 5–9, 10–14, 15–19, 20–24, 25–29, 30–34, 35–39, 40–44, 45–49, 50–54, 55–59, 60–64, 65–69, 70–74, 75+ years), sex (male, female), race (White, Black, mixed, Asian, indigenous), educational level (years of study: 0, 1–3, 4–7, 8–11, 12+, unknown), district of residence, place of death, date of death, type of hospital administration (public state government, public city government, private/non-profit) and cause of death.

We also gathered information on population estimates from SEADE (Statewise System for Data Analysis Foundation) and on the following socio-economic indicators: household crowding (calculated as the average number of persons/household),<sup>14</sup> educational attainment (calculated as the percentage of the population with a university degree),<sup>15</sup> income level (calculated as the percentage of persons living with less than one-quarter of the minimum wage in 2018)<sup>16</sup> and percentage of households located in subnormal areas (irregular settlements, also known as 'favelas').<sup>17</sup> All

variables were collected for each district ( $n = 96$ ) of the city and categorized based on their quintile distribution.

### Analytical methods

Age-standardized mortality rates (ASRs) (2010 Brazilian Census population) and crude rate ratios (RRs) [with corresponding 95% confidence intervals (95% CIs)] were calculated. Mortality rates were expressed as deaths per 100 000 person-years and analysed by race, sex, age group and residential district. Also, to investigate the possible influence of socio-economic factors on COVID-19 mortality, we calculated ASR by household crowding, educational attainment, income level and percentage of subnormal areas in each district. We further split age into two categories ( $<60$  years = young/adult population and  $\geq 60$  years = elderly population), calculating ASR for each group according to race, sex and all socio-economic indicators. We have also evaluated the association between place of death, type of hospital administration and race. We calculated odds ratios (ORs) and 95% CIs. We have tested whether age modifies these associations using the Breslow-Day test for the homogeneity of ORs. These analyses were performed using Stata 15.1 (Stata Corp., College Station, TX, USA).

We also analysed time trends in COVID-19 mortality in São Paulo's city according to race using a Joinpoint Regression Model with the age-standardized mortality rate as the dependent epidemiological week (from Week 12: 15–21 March to Week 39: 20–26 September) as the independent variable. We estimated the weekly percent changes (WPCs) in the mortality rate and corresponding 95% CIs. The number of join points was obtained using a permutation test via Monte Carlo resampling. Once the number  $k$  of join points was obtained, the different models were compared by estimating their Bayesian Information Criterion (BIC).<sup>18</sup> Analyses were performed using the Joinpoint Regression Program version 4.7.0.0 (Joinpoint Regression Program, Version 4.7.0.0. February 2019; Statistical Research and Applications Branch, National Cancer Institute).

### Results

Between March and September 2020, the mortality system recorded 19 498 deaths due to COVID-19, corresponding to 37.1% of all deaths due to diseases that occurred in the city in the same period ( $n = 52 519$ ). Overall, the age-adjusted mortality rate observed in the period was 123.2 deaths/100 000 for all ages.

The highest mortality rate was observed in the district of Brás (192.3 deaths/100 000 inhabitants) and the lowest in Jardim Paulista (48.1 deaths/100 000 inhabitants) (Supplementary Table 1, available as Supplementary data

at *IJE* online). The distribution of the deaths according to age, sex, race, month of death, place of death, ICD-10 code and type of hospital administration is shown in Table 1.

Males presented an 84% increase in mortality due to COVID-19 compared with females (RR = 1.84, 95% CI 1.79–1.90). Analysis by age group revealed that sex differences were similar in the age groups (young/adult population, RR = 2.00, 95% CI 1.88–2.13; elderly population, RR = 1.76, 95% CI 1.70–1.82) (Table 2).

Higher mortality rates were observed for Blacks (RR = 1.77, 95% CI 1.67–1.88) and mixed (RR = 1.42, 95% CI 1.37–1.47) compared with Whites, whereas lower mortality was noted for Asians (RR = 0.63, 95% CI 0.58–0.68). A stratified analysis by age revealed that differences in COVID-19 mortality according to race were more pronounced among persons  $<60$  years old (Blacks, RR = 2.12, 95% CI 1.85–2.42; mixed, RR = 1.62, 95% CI 1.51–1.75; Asians, RR = 0.47, 95% CI 0.38–0.59) than among the elderly (Blacks, RR = 1.68, 95% CI 1.56–1.80; mixed, RR = 1.37, 95% CI 1.31–1.43; Asians, RR = 0.67, 95% CI 0.62–0.73) (Table 2).

A positive gradient was found for all indicators of socio-economic status (SES), i.e. increases in disparities denoted by less education, more household crowding, lower income and a higher concentration of subnormal areas were associated with higher mortality rates. However, a 'dose-response' effect was only observed for education and household density. Among all indicators, the educational level was the one showing the most substantial disparity between the categories. In the young/adult population, among those living in areas with the lowest percentage of the population with a university degree, mortality was four times higher compared with that in the most educated group (RR = 4.02, 95% CI 3.42–4.72); in the elderly population, the same comparison denoted a 96% increased risk of death (Table 2).

Stratified analysis examining racial disparities in mortality according to area-based measures of school attainment revealed that differences between Whites and Blacks were higher in areas with a better educational level. In areas with the lowest percentages of persons with a university degree, Blacks had a risk of death 32% higher than that for Whites, whereas, for those living in the group of districts with the highest percentages of persons with a university degree, this risk was 49% higher than that observed for the White population. Differences between White and mixed populations also persisted in all groups, except in the second and fifth quintiles. Similarly, stratified analysis by household density denoted that higher risk of deaths for Blacks compared with Whites were present in all strata, with relative risks ranging between 1.39 and 1.77;

**Table 1** Distribution of COVID-19 deaths according to age, sex, race school attainment, place of death, month of death, ICD-10 code and hospital administration

Variables	Number of deaths (%)
<b>Age (years)</b>	
0–19	80 (0.5)
20–39	600 (3.8)
40–59	2801 (17.5)
60–74	5163 (32.3)
≥75	7331 (45.9)
<b>Sex</b>	
Male	10 670 (54.7)
Female	8824 (45.3)
Unknown	4 (0.0)
<b>Race</b>	
White	11 946 (61.3)
Black	1793 (9.2)
Asian	444 (2.3)
Mixed	4644 (23.8)
Indigenous	13 (0.1)
Unknown	658 (3.4)
<b>School attainment (years)</b>	
0 (illiterate)	1515 (7.8)
1–7	9413 (48.3)
≥8	6250 (32.0)
Unknown	2320 (11.9)
<b>Place of death</b>	
Hospital	18 253 (93.6)
Other health facilities	447 (2.3)
Home	709 (3.6)
Other places	89 (0.5)
<b>Month of death</b>	
March	511 (2.6)
April	3529 (18.1)
May	5000 (25.7)
June	4061 (20.8)
July	2929 (15.0)
August	2268 (11.6)
September	1200 (6.2)
<b>ICD-10 code</b>	
B34.2 (confirmed)	13 801 (70.8)
U04.9 (suspect)	5697 (29.2)
<b>Type of hospital administration<sup>a</sup></b>	
Public/state government	5144 (29.3)
Public/city government	5196 (29.6)
Private/non-profit	7210 (41.1)

<sup>a</sup>Only for deaths that occurred in hospitals/other health facilities located in São Paulo city ( $n = 17\,550$ ).

<sup>\*\*</sup>Following guidance from the Ministry of Health of March 2020, code U04.9 (Severe acute respiratory syndrome, unspecified) was used as a marker for suspected cases of COVID-19 and code B34.2 (Coronavirus infection, unspecified site) was used for confirmed cases of the disease (SARS-CoV-2 detected by RT-PCR).

accordingly, the same pattern was observed when comparing mixed to White populations, although with increases in the risk of death ranging from 6% to 39% (Table 3).

We observed an association between race and place of death. Whereas Asians were the group with the highest percentage of home deaths (6.8%), Blacks (3.2%) and mixed populations (3.1%) presented the highest percentages of deaths in health facilities other than hospitals, i.e. primary-care units and isolated emergency-care units. Compared with Whites, the Black and mixed populations presented higher risks of dying in other health facilities (67% and 64%, respectively), whereas Asians had a higher probability of dying at home (OR = 1.67, 95% CI 1.14–2.45). The distribution of deaths according to the type of health-facility administration showed that Blacks and mixed individuals died predominantly in public institutions (75.5% and 76.5%, respectively). In comparison, deaths among Whites (49.4%) and Asians (72.2%) were concentrated in private or non-profit health facilities. We observed that age modifies this association, with larger effects being observed for the groups of elderly Blacks and mixed (Table 4).

In the White population, we observed a significant increase in COVID-19 mortality between Weeks 12 and 14 (WPC = 158.8%, 95% CI 37.7–386.4), followed by another increase between epidemiological Weeks 14 and 21 (WPC = 9.6%, 95% CI 5.3–14.0) and a decrease between Weeks 21 and 39 (WPC = -7.4%, 95% CI -8.3 to -6.5). Overall, for the entire period, there was an average weekly increase of 4.4% in COVID-19 mortality in this group (AWPC = 4.4, 95% CI -0.3 to 9.2). Among Blacks, a non-significant increase in COVID-19 mortality between Weeks 12 and 14 was also noted (WPC = 377.5%, 95% CI -34.7 to 3392.4), followed by an increase between epidemiological Weeks 14 and 23 (WPC = 7.3%, 95% CI 2.5–12.3) and a decrease between Weeks 23 and 39 (WPC = -9.6%, 95% CI -11.4 to -7.8). Overall, there was an average weekly increase of 16.8% in COVID-19 mortality in this group (AWPC = 8.3, 95% CI -5.9 to 24.5). For the mixed group, the model identified more join points, with a large increase in COVID-19 mortality between Weeks 12 and 14 (WPC = 309.0%, 95% CI 13.3–1376.9), followed by another increase between epidemiological Weeks 14 and 19 (WPC = 18.1%, 95% CI 9.3–27.5) and decreases in rates between Weeks 19 and 24 (WPC = -3.2%, 95% CI -9.6 to 3.6), Weeks 24 and 27 (WPC = -16.6%, 95% CI -35.8 to 8.5), Weeks 27 and 32 (WPC = -1.8%, 95% CI -10.8 to +8.1) and Weeks 32 and 39 (WPC = -13.2%, 95% CI -17.9 to -8.3); the average weekly percentage change in this population was 7.1% (95% CI -2.3 to 17.5). Asians showed the lowest increase in mortality in the study period (WPC 1: Weeks 12–15 = 94.0%, 95% CI -0.1 to 276.2; WPC 2: Weeks 15–39 = -4.6%, 95% CI -7.4 to -1.7; AWPC = 6.5%, 95% CI -4.2 to 18.3) (Figure 1).

**Table 2** Number of deaths, age-standardized mortality rates (/100 000 hab) and rate ratios (95% CI) of COVID-19 according to education attainment, sex, race, household density, household income, percentage of households in subnormal areas and age group, São Paulo city, March–September 2020

Variables	Age group											
	<60						≥60					
	n	ASR	RR	95% CI	n	ASR	RR	95% CI	n	ASR	RR	95% CI
<b>Population with university degree (%)</b>												
<8.61	5301	151.6	2.23	2.13–2.34	1477	40.2	4.02	3.42–4.72	3824	1072.3	1.96	1.86–2.06
8.61–13.20	4812	130.8	1.92	1.83–2.02	1127	34.5	3.45	2.92–4.08	3685	926.5	1.69	1.61–1.78
13.21–19.20	3308	122.2	1.73	1.65–1.82	610	32.5	3.25	2.75–3.84	2698	863.2	1.58	1.49–1.66
19.21–34.80	2940	101.8	1.57	1.48–1.67	436	22.7	2.27	1.89–2.72	2504	755.3	1.38	1.31–1.46
>34.80	2277	68.0	Ref.	Ref.	169	10.0	Ref.	Ref.	2108	547.7	Ref.	Ref.
<b>Sex</b>												
Male	10 670	173.4	1.84	1.79–1.90	2579	72.2	2.00	1.88–2.13	8086	1713.3	1.76	1.70–1.82
Female	8824	94.2	Ref.	Ref.	1494	36.1	Ref.	Ref.	7329	974.3	Ref.	Ref.
<b>Race<sup>a</sup></b>												
White	11 946	106.2	Ref.	Ref.	1979	25.5	Ref.	Ref.	9964	772.7	Ref.	Ref.
Black	1793	188.1	1.77	1.67–1.88	475	54.0	2.12	1.85–2.42	1317	1296.2	1.68	1.56–1.80
Asian	444	66.7	0.63	0.58–0.68	42	12.0	0.47	0.38–0.59	402	518.5	0.67	0.62–0.73
Mixed	4644	150.8	1.42	1.37–1.47	1416	41.4	1.62	1.51–1.75	3226	1055.2	1.37	1.31–1.43
Indigenous	13	96.2	0.91	0.54–1.52	5	46.2	1.81	0.56–5.81	8	509.7	0.66	0.38–1.15
<b>Mean household density (persons/room)</b>												
<2.71	2340	69.9	Ref.	Ref.	182	10.4	Ref.	Ref.	2158	561.3	Ref.	Ref.
2.71–2.88	3720	107.7	1.54	1.46–1.62	618	26.5	2.55	2.18–2.97	3102	778.7	1.39	1.31–1.47
2.89–2.99	3918	112.0	1.60	1.52–1.69	845	30.8	2.96	2.57–3.41	3073	857.2	1.53	1.45–1.61
3.00–3.07	4235	141.5	2.02	1.92–2.13	994	38.1	3.66	3.21–4.18	3241	995.7	1.77	1.68–1.87
>3.07	4425	146.8	2.10	2.00–2.21	1180	39.4	3.79	3.55–4.28	3245	1034.2	1.84	1.74–1.95
<b>Population with monthly income &lt;1/4 minimum wage (%)</b>												
<5.34	2780	88.6	Ref.	Ref.	330	17.2	Ref.	Ref.	2450	678.9	Ref.	Ref.
5.34–6.77	3456	107.1	1.21	1.15–1.27	574	26.8	1.56	1.37–1.77	2882	770.9	1.14	1.08–1.20
6.78–8.28	2838	99.3	1.12	1.06–1.18	466	24.7	1.44	1.26–1.64	2372	715.6	1.05	1.00–1.12
8.29–10.65	4965	133.8	1.51	1.44–1.58	1131	34.1	1.98	1.78–2.21	3834	938.2	1.41	1.34–1.48
>10.65	4599	149.6	1.69	1.61–1.77	1318	40.4	2.35	2.12–2.60	3281	1052.0	1.55	1.47–1.63
<b>Percentage of households in subnormal areas (%)</b>												
<0.65	2501	83.1	Ref.	Ref.	274	16.5	Ref.	Ref.	2227	633.5	Ref.	Ref.
0.65–3.05	2579	100.0	1.20	1.14–1.27	388	25.2	1.53	1.31–1.79	2191	718.2	1.13	1.07–1.20
3.06–6.67	4249	129.2	1.55	1.48–1.63	915	35.0	2.12	1.87–2.40	3334	907.9	1.43	1.36–1.51
6.68–12.56	3918	138.7	1.67	1.59–1.76	970	37.9	2.30	2.04–2.59	2948	971.9	1.45	1.37–1.51
>12.56	5391	126.9	1.53	1.46–1.60	1272	31.7	1.92	1.72–2.15	4119	913.4	1.44	1.37–1.52

<sup>a</sup>Deaths with unknown age = 3 White, 1 Black, 2 mixed, 1 unknown race. ASR, age-standardized rate; RR, rate ratio.

**Table 3** Number of deaths, age-standardized mortality rates (/100,000 hab) and rate ratios (95% CI) of COVID-19 according to race, stratified by education attainment and household density, São Paulo city, March–September 2020

Variables	Race																			
	Quintiles			White			Black			Asian			Mixed			Indigenous				
	n	ASR	RR	n	ASR	RR	95% CI	n	ASR	RR	95% CI	n	ASR	RR	95% CI	n	ASR	RR	95% CI	
Percentage of the population with university degree	<8.61	2508	139.2	Ref.	647	183.3	1.32	1.20-1.45	24	58.9	0.42	0.33-0.55	1913	147.5	1.06	1.00-1.13	6	129.1	0.93	0.43-2.01
	8.61-13.20	2708	119.3	Ref.	537	186.5	1.56	1.40-1.74	60	99.1	0.83	0.66-1.05	1365	110.9	0.93	0.87-0.99	5	108.8	0.91	0.39-2.11
	13.21-19.20	2182	106.7	Ref.	275	187.4	1.75	1.49-2.04	110	102.0	0.96	0.79-1.15	643	117.4	1.10	1.01-1.20	1	58.8	0.55	0.13-2.31
	19.21-34.80	2121	91.5	Ref.	208	178.5	1.95	1.61-2.36	115	65.1	0.71	0.60-0.84	395	125.8	1.37	1.22-1.55	1	38.4	0.42	0.12-1.44
	>34.80	1909	64.4	Ref.	51	96.1	1.49	1.07-2.09	124	56.9	0.88	0.74-1.05	98	69.4	1.08	0.87-1.33	0	0.0	-	-
Household density (persons/household)	<2.71	1931	65.2	Ref.	64	111.5	1.71	1.24-2.35	129	59.9	0.92	0.77-1.09	119	77.5	1.19	0.97-1.45	0	0.00	-	-
	2.71-2.88	2648	97.1	Ref.	256	172.0	1.77	1.50-2.09	129	67.5	0.70	0.60-0.81	583	135.3	1.39	1.26-1.54	1	46.5	0.48	0.13-1.81
	2.89-2.99	2283	106.6	Ref.	373	163.0	1.53	1.35-1.74	83	74.1	0.70	0.58-0.84	1030	133.3	1.25	1.16-1.35	2	53.9	0.51	0.19-1.33
	3.00-3.07	2341	141.9	Ref.	511	209.6	1.48	1.32-1.65	62	88.7	0.63	0.51-0.76	1183	156.5	1.10	1.03-1.18	3	104.4	0.74	0.28-1.94
>3.07	2225	163.6	Ref.	514	226.7	1.39	1.24-1.54	30	69.4	0.42	0.34-0.54	1499	172.8	1.06	0.99-1.13	7	195.6	1.20	0.53-2.69	

ASR, age-standardized rate; RR, rate ratio.

## Discussion

To our knowledge, this is the first population-based study addressing racial disparities and examining time trends in COVID-19 mortality in the most populous city in Latin America. We showed that Blacks and mixed individuals observed mortality rates 81% and 45% higher than Whites, respectively, and disparities were more pronounced in the young/adult population. It is also noteworthy that Asians presented a decreased risk of death compared with Whites. Besides, the analysis of mortality rates by other indicators such as income, education and housing revealed a social gradient, i.e. worsening in the indicators increases mortality rates. Our results corroborate previous findings indicating sex, race and socioeconomic disparities in COVID-19 mortality worldwide.<sup>19-21</sup>

A previous Brazilian study using data from 11 321 patients admitted to hospitals included in the SIVEP-Gripe data set (Influenza Epidemiological Surveillance System) has shown that mixed and Blacks with COVID-19 presented a higher risk of death compared with Whites (45% and 32%, respectively), with disparities likely explained by differences in exposure to infection by SARS-Cov-2 and access to healthcare.<sup>5</sup> In an ecological study, Martins-Filho *et al.* found that, in Aracaju, the capital of Sergipe state (Northeast region), case-fatality rates increased concomitantly with decreasing values of a living-condition index (LCI) (including area-based information on education, income and housing); whereas the case-fatality rate was 0.8% in the neighbourhoods with high LCI, those areas with very low LCI had a value of 2%.<sup>22</sup> Also, Santos *et al.* described that, in Brazil, maternal mortality due to COVID-19 among Blacks was almost twice the value observed for White women. Although Black and White women were similar in age and co-morbidities burden, they differed in the clinical status at hospital admission, with Black women being hospitalized in worse conditions, suggesting that these disparities could be linked to a context outside of the hospitals.<sup>23</sup>

Using data from 27 states in the USA and the city of New York, Gross *et al.* reported a strong association between Black race, Hispanic ethnicity and COVID-19 mortality. Overall, the mortality rate among blacks was 3.6 times higher than that among Whites, whereas Hispanics presented a 88% increase in mortality.<sup>24</sup> Results from a national survey conducted in the USA demonstrated that, compared with Whites, African Americans, younger individuals and men reported more COVID-19 exposure, less social isolation and had less detailed knowledge about the disease.<sup>25</sup>

**Table 4** Number and percentage of COVID-19 deaths and odds ratios according to race, place of death, type of hospital administration and age group, São Paulo city, March–September, 2020

Race	<60											≥60												
	All ages		Other health facilities (OHF)		Home (HO)		Crude OR H vs OHF (95% CI)		Hospital (H)		Other health facilities (OHF)		Home (HO)		Crude OR H vs OHF (95% CI)		Hospital (H)		Other health facilities (OHF)		Home (HO)		Crude OR H vs OHF (95% CI)	
White	11 170 (94.0)	227 (1.9)	491 (4.1)	Ref. (1.9)	1863 (94.8)	35 (1.8)	68 (3.4)	Ref. (3.4)	9305 (93.8)	192 (1.9)	422 (4.3)	Ref. (4.3)	9305 (93.8)	192 (1.9)	422 (4.3)	Ref. (4.3)	9305 (93.8)	192 (1.9)	422 (4.3)	Ref. (4.3)	422 (4.3)	Ref. (4.3)	Ref. (4.3)	Ref. (4.3)
Black	1676 (93.8)	57 (3.2)	53 (3.0)	1.67 (1.25–2.25)	445 (94.1)	13 (2.7)	15 (3.2)	1.56 (0.82–2.96)	1230 (93.8)	44 (3.3)	38 (2.9)	1.73 (1.24–2.42)	1230 (93.8)	44 (3.3)	38 (2.9)	1.73 (1.24–2.42)	1230 (93.8)	44 (3.3)	38 (2.9)	1.73 (1.24–2.42)	38 (2.9)	1.73 (1.24–2.42)	0.68 (0.49–0.95)	
Asian	408 (92.5)	3 (0.7)	30 (6.8)	0.36 (0.11–1.13)	39 (92.9)	0 (0.0)	3 (7.1)	NC–	369 (92.5)	3 (0.7)	27 (6.8)	0.39 (0.12–1.24)	369 (92.5)	3 (0.7)	27 (6.8)	0.39 (0.12–1.24)	369 (92.5)	3 (0.7)	27 (6.8)	0.39 (0.12–1.24)	27 (6.8)	0.39 (0.12–1.24)	1.61 (1.08–2.41)	
Mixed	4353 (94.1)	145 (3.1)	127 (2.8)	1.64 (1.33–2.02)	1311 (93.4)	49 (3.5)	44 (3.1)	1.99 (1.28–3.09)	3040 (94.4)	96 (3.0)	83 (2.6)	1.53 (1.19–1.96)	3040 (94.4)	96 (3.0)	83 (2.6)	1.53 (1.19–1.96)	3040 (94.4)	96 (3.0)	83 (2.6)	1.53 (1.19–1.96)	83 (2.6)	1.53 (1.19–1.96)	0.60 (0.47–0.76)	

Type of hospital administration <sup>b</sup> (%)															
Race	Public/ City Gov. (PCG)			Private (P)			Public/ State Gov. (PSG)			Public/ City Gov. (PCG)					
	State Gov. (PSG)	Private (P)	Public/ City Gov. (PCG)	State Gov. (PSG)	Private (P)	Public/ City Gov. (PCG)	State Gov. (PSG)	Private (P)	Public/ City Gov. (PCG)	State Gov. (PSG)	Private (P)	Public/ City Gov. (PCG)			
White	2845 (26.5)	2587 (24.1)	5301 (49.4)	Ref. (31.0)	540 (31.0)	541 (31.0)	Ref. (38.0)	662 (38.0)	540 (31.0)	541 (31.0)	Ref. (51.6)	2306 (25.6)	2046 (22.8)	4636 (51.6)	
Black	586 (36.4)	631 (39.1)	395 (24.5)	2.76 (2.41–3.16)	144 (34.7)	169 (40.7)	1.73 (1.31–2.28)	102 (24.6)	144 (34.7)	169 (40.7)	2.03 (1.55–2.66)	442 (37.0)	461 (38.5)	293 (24.5)	3.03 (2.59–3.54)
Asian	64 (16.2)	46 (11.6)	286 (72.2)	0.42 (0.32–0.55)	8 (22.2)	7 (19.4)	0.47 (0.20–1.06)	21 (58.4)	8 (22.2)	7 (19.4)	0.41 (0.17–0.97)	56 (15.6)	39 (10.8)	265 (73.6)	0.42 (0.32–0.57)
Mixed	1486 (35.5)	1713 (41.0)	984 (23.5)	2.81 (2.56–3.09)	412 (33.0)	524 (41.9)	1.61 (1.34–1.94)	314 (25.1)	412 (33.0)	524 (41.9)	2.04 (1.70–2.44)	1073 (36.6)	1188 (40.5)	670 (22.9)	3.22 (2.89–3.59)

OR, odds ratio.

<sup>a</sup>Excludes deaths that occurred in other places and deaths with indigenous or unknown race ( $n = 758$ ).

<sup>b</sup>Includes only deaths that occurred in São Paulo city in hospitals or other facilities ( $n = 17 550$ ); excludes deaths with indigenous or unknown race ( $n = 638$ ).


Breslow-Day test for interactions between race and age.

Place of death (hospital vs other health facilities): Blacks = chi-square = 0.089,  $P = 0.769$ ; Asians = not calculated; Mixed = chi-square = 1.037,  $P = 0.310$ .

Place of death (hospital vs home): Blacks = chi-square = 0.814,  $P = 0.367$ ; Asians = chi-square = 0.171,  $P = 0.679$ ; Mixed = chi-square = 3.348,  $P = 0.067$ .

Type of hospital administration (private vs public state government): Blacks = chi-square = 11.9,  $P = 0.001$ ; Asians = chi-square = 0.045,  $P = 0.832$ ; Mixed = chi-square = 39.97,  $P < 0.001$ .

Type of hospital administration (private vs public city government): Blacks = chi-square = 12.56,  $P < 0.001$ ; Asians = chi-square = 0.181,  $P = 0.670$ ; Mixed = chi-square = 39.86,  $P < 0.001$ .



**Figure 1** Trends in COVID-19 mortality according to epidemiological week and race, São Paulo city, March–July 2020

Our results show that COVID-19 mortality was 84% higher among males than that among females, with an effect persisting in both young/adults and elderly groups. This finding is consistent with several previous reports showing an excess of COVID-19 mortality in men.<sup>26–28</sup> It is still debatable whether these disparities are attributable to biologic factors such as sex hormones and expression of ACE2 receptor,<sup>29</sup> differences in lifestyle that could result in men having more co-morbidities,<sup>30–31</sup> differences in the exposure to SARS-CoV-2 due to behavioural and occupational factors<sup>29</sup> or symptoms awareness,<sup>32</sup> and differences in adherence to preventive health measures.<sup>33</sup> Elucidating whether the mechanisms behind these differences are linked to social or biological factors is crucial to designing better strategies for COVID-19 prevention and treatment.<sup>34</sup>

Differences in COVID-19 mortality according to education, income and housing conditions described in our study are concordant with findings from a previous ecological study that has used a composite socio-economic index and spatio-temporal analyses to show that socio-economic conditions and COVID-19 mortality were associated in the city of São Paulo.<sup>6</sup> In the UK, COVID-19 mortality rates in the most deprived areas were almost 100% higher than

those observed in the least deprived regions, and the effect was not modified after controlling for ethnicity. Similarly to our study, authors reported that, after adjustment for age, geographic region, sex and ethnicity, social disparities were more pronounced among people of working age (most deprived compared with least deprived, 93% higher risk of death) than among elderly persons (most deprived compared with least deprived, 9% higher risk of death).<sup>10</sup> Evidence suggests that social disparities in COVID-19 mortality may reflect a combination of factors resulting in an increased vulnerability to being exposed to the virus, such as household crowding, underlying medical conditions, occupation and modes of transportation.<sup>35</sup>

We found a higher difference between mortality rates among Blacks and Whites in more affluent areas (higher education and lower household density). According to Telles, residential segregation by race in Brazil is lowest for the low-income groups and increases with increasing income level.<sup>36</sup> We hypothesized that our findings are related to inequalities in job opportunities, as Blacks are disproportionately overrepresented as domestic employees,<sup>37</sup> with these deaths denoting people working and living in the same place. Data from the 2015 National Household

Sample Survey (PNAD) show that, in the Southeast region, 0.7% of all mixed and Black females working as domestic employees lived in the same household in which they worked, but only 52.2% were legally registered; this may reflect lower income and difficulties in immediate access to health services.<sup>38</sup>

Our finding of a higher percentage of deaths occurring in primary-care or isolated emergency-care units observed for Black and mixed populations may be related to barriers in healthcare access. Racial disparities in medical care, both quantitatively and qualitatively, are well documented, being described as a possible contributor to racial disparities in health outcomes.<sup>39</sup> The city of São Paulo has 4.09 intensive-care-unit (ICU) beds/10 000 inhabitants, but this rate is 5.27 in the private sector and only 1.58 ICU beds/10 000 inhabitants for the public one [Sistema Único de Saúde (SUS)].<sup>40</sup> Data from SIVEP-Gripe for the same period of our study show that Asians (39.0%) and Whites (35.7%) were more frequently admitted to ICUs than Blacks (29.8%), mixed (30.4%) and indigenous (24.7%). One could argue that differences in disease severity could explain this, but an analysis of a very high-risk group (persons with diabetes, O<sub>2</sub> saturation <95% and symptoms of respiratory distress) also revealed these disparities in ICU admission (Whites = 46.2%, Blacks = 40.2%, Asians = 47.0%, mixed = 39.9%).<sup>41</sup> In New York City, which has an extensive hospital system, a study assessing racial-area composition based on zip codes has demonstrated a striking disparity in hospital resources during pandemics, with predominantly White areas having almost nine times more beds/1000 persons than predominantly Black/Hispanic areas.<sup>42</sup>

Despite being the group with the lowest COVID-19 mortality, a higher proportion of home deaths was found among Asians. Further exploration of our data shows that this effect was even more substantial when we restricted the analysis to COVID-19 suspected deaths. This finding could be explained by the fear of seeking medical care in the context of the pandemics (misinterpreting the symptoms) or cultural differences/preferences on facing a severe disease and the risk of death, particularly among elderly persons.<sup>43</sup>

Time-trend analyses showed an earlier decrease in mortality among Asians and Whites compared with that among Blacks and mixed. Possible explanations include sustained higher incidence and mortality rates among blacks and mixed due to difficulties in adhering to social-distancing measures. Coelho *et al.* reported the challenges of implementing social-distancing and adequate hygiene measures in areas with high social vulnerability due to household crowding and poor living conditions.<sup>44</sup> In the USA, the disparate impact of COVID-19 in the Latino community has been linked to work and living conditions and financial burden, as paid sick leave and working from

home are not options for many Hispanic workers.<sup>7</sup> In Brazil, the situation is not different. Pires *et al.* reported that we should expect that more vulnerable Brazilians will be disproportionately affected by COVID-19, not only because of the previously cited reasons, but also because there is a high incidence of co-morbidities among persons with less education.<sup>45</sup>

Limitations of our study include the use of area-based SES measures for relatively large geographical units (districts) and the lack of detailed information about co-morbidities. Regarding ‘ecological fallacy’, it is important to emphasize that we are aware of this bias and the limitations of using area-based measures for relatively large geographic units (districts). However, data from the literature suggest that heterogeneity in the neighbourhood would probably produce underestimation and not overestimation of the effects.<sup>46–47</sup> Also, it is essential to mention the possibility of racial misclassification, as, in Brazil, this is self-reported mostly based on skin colour and it is strongly affected by the context and social status.<sup>48</sup> The strengths of the study lie in the quality of the mortality data and its completeness. In the state of São Paulo, the coverage and completeness of death registration are equal to 100%<sup>49</sup> and only 11.4% of the deaths were reported as ‘garbage codes’ (unpublished observation).

In conclusion, our results show social inequalities in COVID-19 mortality in São Paulo, Brazil. Unfortunately, the relationship between more vulnerable = less health is not a new phenomenon, and we believe that pandemics have exacerbated existing structural inequalities. Stiglitz correctly states that ‘COVID-19 is not an equal opportunity virus’, emphasizing that differences in the success (or not) in the management of the pandemics in each country are linked to several factors including the healthcare system, pre-existing health inequalities, economy’s resilience, reliance on science and citizens’ trust in government guidance.<sup>50</sup> In Brazil, the governmental response to COVID-19 has been marked by the lack of leadership at the federal level, distrust of science, denial of the importance of the virus and progressive cuts to health and research funding. ‘So what?’<sup>51–52</sup> We want politicians to remember what is written in our Constitution: ‘Health is a right for all Brazilian citizens and a duty of the State, guaranteed through social and economic policies designed to reduce the risk of disease and promote universal and equal access to healthcare.’<sup>53</sup> Until equity and trust in science become essential elements in the Brazilian response, COVID-19 will continue to highlight the sorrowful shades of social disparities in the largest city of Latin America.

## Supplementary Data

Supplementary data are available at *IJE* online.

## Funding

None.

## Acknowledgements

Data derived from a source in the public domain (Sistema de Informações sobre Mortalidade—SIM/PRO-AIM/CEInfo-SMS/SP) available at [http://tabnet.saude.prefeitura.sp.gov.br/cgi/deftohtm3.exe?secretarias/saude/TABNET/SIM\\_PROV/obitop.def](http://tabnet.saude.prefeitura.sp.gov.br/cgi/deftohtm3.exe?secretarias/saude/TABNET/SIM_PROV/obitop.def). Ethics approval was not necessary because this study analysed only mortality publicly available data, not including identifiable information.

## Conflict of Interest

None declared.

## References

- Zhu N, Zhang D, Wang W *et al*. A novel coronavirus from patients with pneumonia in China, 2019. *N Engl J Med* 2020; 382:727–33.
- World Health Organization. *Coronavirus Disease 2019 (COVID-19)—Situation Report 51*. Geneva: World Health Organization.
- World Health Organization. WHO Coronavirus Disease (COVID-19) Dashboard. <https://covid19.who.int/> (12 October 2020, date last accessed).
- Ministério da Saúde. COVID-19 NO BRASIL. <https://covid.saude.gov.br/> (14 October 2020, date last accessed).
- Baqui P, Bica I, Marra V, Ercole A, van der Schaar M. Ethnic and regional variations in hospital mortality from COVID-19 in Brazil: a cross-sectional observational study. *Lancet Glob Health* 2020;8:e1018–26.
- Bermudi PMM, Lorenz C, Aguiar BS, Failla MA, Barrozo LV, Chiaravalloti-Neto F. Spatiotemporal ecological study of COVID-19 mortality in the city of São Paulo, Brazil: shifting of the high mortality risk from areas with the best to those with the worst socio-economic conditions. *Travel Med Infect Dis* 2021; 39:101945.
- Macias Gil R, Marcelin JR, Zuniga-Blanco B, Marquez C, Mathew T, Piggott DA. COVID-19 pandemic: disparate health impact on the Hispanic/Latinx population in the United States. *Journal of Infectious Diseases* 2020;222:1592–95.
- Baidal JW, Wang AY, Zumwalt K *et al*. Social determinants of health and COVID-19 among patients in New York City. *Res Sq* 2020; doi: 10.21203/rs.3.rs-70959/v1.
- Dorn AV, Cooney RE, Sabin ML. COVID-19 exacerbating inequalities in the US. *Lancet* 2020;395:1243–44.
- Public Health England. *Disparities in the Risk and Outcomes of COVID-19*. London. <https://www.gov.uk/government/publications/covid-19-review-of-disparities-in-risks-and-outcomes> (5 October 2020, date last accessed).
- Amengual-Moreno M, Calafat-Caules M, Carot A *et al*. [Social determinants of the incidence of Covid-19 in Barcelona: a preliminary ecological study using public data.]. *Rev Esp Salud Publica* 2020;94:e202009101.
- Góes C, Karpowicz I. *Inequality in Brazil: A Regional Perspective*, International Monetary Fund Working Papers 2017. <https://www.imf.org/en/Publications/WP/Issues/2017/10/31/Inequality-in-Brazil-A-Regional-Perspective-45331> (10 October 2020, date last accessed).
- PRO-AIM/CEINFO-SMS-SP. *Sistema de Informações sobre Mortalidade*. <https://www.prefeitura.sp.gov.br/cidade/secretarias/saude/tabnet/mortalidade/index.php?p=6529> (5 October 2020, date last accessed).
- SEADE—Fundação Sistema Estadual de Análise de Dados. *Sistema SEADE de Projeções Populacionais*. <http://produtos.seade.gov.br/produtos/projpop/> (2 October 2020, date last accessed).
- Companhia do Metropolitan de São Paulo. *Pesquisa Origem Destino—2017*. <http://www.metro.sp.gov.br/pesquisa-od/> (30 September 2020, date last accessed).
- SEADE—Fundação Sistema Estadual de Análise de Dados. *Informações dos municípios paulistas*. <http://www.imp.seade.gov.br/frontend/#/> (2 October 2020, date last accessed).
- Secretaria Municipal de Desenvolvimento Urbano de São Paulo. Observatório de Indicadores da Cidade de São Paulo. <http://observampa.prefeitura.sp.gov.br/> (30 September 2020, date last accessed).
- Kim HJ, Fay MP, Feuer EJ, Midthune DN. Permutation tests for joinpoint regression with applications to cancer rates. *Statist Med* 2000;19:335–51.
- Ahrenfeldt LJ, Otavova M, Christensen K, Lindahl-Jacobsen R. Sex and age differences in COVID-19 mortality in Europe. *Res Sq* 2020; doi: 10.21203/rs.3.rs-61444/v1, preprint: not peer-reviewed.
- Golestaneh L, Neugarten J, Fisher M *et al*. The association of race and COVID-19 mortality. *E Clinical Medicine* 2020;25: 100455.
- Williamson EJ, Walker AJ, Bhaskaran K *et al*. Factors associated with COVID-19-related death using OpenSAFELY. *Nature* 2020;584:430–36.
- Martins-Filho PR, de Souza Araujo AA, Quintans-Junior LJ, Santos VS. COVID-19 fatality rates related to social inequality in Northeast Brazil: a neighborhood-level analysis. *J Travel Med* 2020;27:taaa128.
- Santos DS, Menezes MO, Andreucci CB *et al*. Disproportionate impact of COVID-19 among pregnant and postpartum Black Women in Brazil through structural racism lens. *Clin Infect Dis* 2020. Published online 2020 Jul 28. doi: 10.1093/cid/ciaa1066
- Gross CP, Essien UR, Pasha S, Gross JR, Wang SY, Nunez-Smith M. Racial and ethnic disparities in population-level Covid-19 mortality. *J Gen Intern Med* 2020;35:3097–99.
- Alsan M, Stantcheva S, Yang D, Cutler D. Disparities in coronavirus 2019 reported incidence, knowledge, and behavior among US adults. *JAMA Netw Open* 2020;3:e2012403.
- Chen T, Wu D, Chen H *et al*. Clinical characteristics of 113 deceased patients with coronavirus disease 2019: retrospective study. *BMJ* 2020;368:m1091.
- Li LQ, Huang T, Wang YQ *et al*. COVID-19 patients' clinical characteristics, discharge rate, and fatality rate of meta-analysis. *J Med Virol* 2020;92:577–83.
- Jin JM, Bai P, He W *et al*. Gender differences in patients with COVID-19: focus on severity and mortality. *Front Public Health* 2020;8:152.
- Kopel J, Perisetti A, Roghani A, Aziz M, Gajendran M, Goyal H. Racial and gender-based differences in COVID-19. *Front Public Health* 2020;8:418.

30. Elgendy IY, Pepine CJ. Why are women better protected from COVID-19: clues for men? Sex and COVID-19. *Int J Cardiol* 2020;315:105–6.
31. Burstrom B, Tao W. Social determinants of health and inequalities in COVID-19. *Eur J Public Health* 2020;30:617–18.
32. Sun Z, Yang B, Zhang R, Cheng X. Influencing factors of understanding COVID-19 risks and coping behaviors among the elderly population. *Ijerp* 2020;17:5889.
33. Okten IG, Oettingen G. Gender differences in preventing the spread of coronavirus. *PsyArXiv* 2020; doi: 10.31234/osf.io/ch4jy, preprint: not peer-reviewed.
34. Islam N, Khunti K, Dambha-Miller H, Kawachi I, Marmot M. COVID-19 mortality: a complex interplay of sex, gender, and ethnicity. *Eur J Public Health* 2020;30:847–48.
35. Blundell R, Costa Dias M, Joyce R, Xu X. COVID-19 and inequalities. *Fisc Stud* 2020; doi: 10.1111/1475-5890.12232 [Online ahead of print].
36. Telles EE. Residential segregation by skin color in Brazil. *Am Sociological Rev* 1992;57:186–97.
37. Lovell PA. Race, gender and regional labor market inequalities in Brazil. *Rev Social Econ* 2000;58:277–93.
38. Instituto de Pesquisa Econômica Aplicada. *Retrato das Desigualdades de Gênero e Raça*. <https://www.ipea.gov.br/retrato/apresentacao.html> (23 October 2020, date last accessed).
39. Williams DR, Rucker TD. Understanding and addressing racial disparities in health care. *Health Care Financ Rev* 2000;21:75–90.
40. Conselho Federal de Medicina. *Menos de 10% dos municípios brasileiros possuem leito de UTI*. [https://portal.cfm.org.br/index.php?option=com\\_content&view=article&cid=27828:2018-09-04-19-31-41&catid=3](https://portal.cfm.org.br/index.php?option=com_content&view=article&cid=27828:2018-09-04-19-31-41&catid=3) (2 October 2020, date last accessed).
41. Secretaria Municipal de Saúde de São Paulo. SIVEP-Gripe. São Paulo. <http://tabnet.saude.prefeitura.sp.gov.br/cgi/deftohtm3.exe?secretarias/saude/TABNET/RSRAG/sragh.def> (23 October 2020, date last accessed).
42. Douglas JA, Subica AM. COVID-19 treatment resource disparities and social disadvantage in New York City. *Prev Med* 2020; 141:106282.
43. Leite AKF, Ribeiro KB. Older adults with cancer in the city of Sao Paulo: what factors determine the place of death? *Rev Saúde Pública* 2018;52:66.
44. Coelho FC, Lana RM, Cruz OG *et al*. Assessing the spread of COVID-19 in Brazil: Mobility, morbidity and social vulnerability. *PLoS One* 2020;15:e0238214.
45. Pires LN, Carvalho L, Xavier LL. *COVID-19 e Desigualdade no Brasil*. CEBES. <http://cebes.org.br/2020/04/covid-19-e-desigualdade-no-brasil/> (20 October 2020, date last accessed).
46. Hyndman JC, Holman CD, Hockey RL, Donovan RJ, Corti B, Rivera J. Misclassification of social disadvantage based on geographical areas: comparison of postcode and collector's district analyses. *Int J Epidemiol* 1995;24:165–76.
47. Krieger N. Women and social class: a methodological study comparing individual, household, and census measures as predictors of Black/White differences in reproductive history. *J Epidemiol Community Health* 1991;45:35–42.
48. Miranda M. *Classificação de Raça/Cor e Etnia: conceitos, Terminologia e Métodos Utilizados Nas Ciências da Saúde No Brasil*. Thesis. Escola Nacional de Saúde Pública, Rio de Janeiro 2010.
49. Queiroz BL, Freire F, Gonzaga MR, Lima EEC. Completeness of death-count coverage and adult mortality (45q15) for Brazilian states from 1980 to 2010. *Rev Bras Epidemiol* 2017; 20:21–33.
50. Stiglitz J. Conquering the Great Divide. *Finance & Development*. 2020 September. International Monetary Fund. <https://www.imf.org/external/pubs/ft/fandd/2020/09/COVID19-and-global-inequality-joseph-stiglitz.htm> (20 October 2020, date last accessed).
51. The Lancet. COVID-19 in Brazil: 'so what?'. *Lancet* 2020;395: 1461.
52. Barberia LG, Gómez EJ. Political and institutional perils of Brazil's COVID-19 crisis. *Lancet* 2020;396:367–8.
53. Presidency of the Republic - Civil House. Constitution of the federative Republic of Brazil of 1988. In: *Presidency of the Republic—Civil House*, editor. Brasília-DF; 1988.